

投与前の血清 中総IgE濃度 (IU/mL)	体重 (kg)									
	≥20~25	>25~30	>30~40	>40~50	>50~60	>60~70	>70~80	>80~90	>90~125	>125~150
≥30~100										
>100~200										
>200~300										375mg
>300~400									450mg	525mg
>400~500							375mg	375mg	525mg	600mg
>500~600						375mg	450mg	450mg	600mg	
>600~700		225mg			375mg	450mg	450mg	525mg		
>700~800	225mg	225mg	300mg	375mg	450mg	450mg	525mg	600mg		
>800~900	225mg	225mg	300mg	375mg	450mg	525mg	600mg			
>900~1,000	225mg	300mg	375mg	450mg	525mg	600mg				
>1,000~1,100	225mg	300mg	375mg	450mg	600mg					
>1,100~1,200	300mg	300mg	450mg	525mg	600mg					投与不可
>1,200~1,300	300mg	375mg	450mg	525mg						
>1,300~1,500	300mg	375mg	525mg	600mg						

2週間毎投与の表に該当しない場合には
4週間毎投与の表に従い投与すること


0.008mg/kg/ IU/mL 2 0.016mg/kg/ IU/mL
4

PD

IgE

IgE

Image



2. IV 2210¹⁾

1) Zielen, S. et al.: Int. Arch. Allergy Immunol. 160(1), 102, 2013 XOLM01866

5

Image



2 ABP EAR

8 EAR 23.1% Group1 9.3% Group2 5.6% p=0.018 p<0.001 ANCOVA 16 EAR 20.0% Group1 11.8% Group2 5.1% p=0.087 p<0.001 ANCOVA

4.


5.

IgE

- FEV1.0 80%
-
- 1

3 PK

Image



4

Group1 50.0% 9/18 Group2 100.0% 16/16 75.0% 12/16
Group1 3 16.7% Group2 5 31.3% 4 25.0%
Group1 1 5.6% Group2 1 6.3% 2.9% 1/34 6.3% 1/16

PFS

IC

IC

IgE

Image



Source URL: <https://prod.pro.novartis.com/jp-ja/products/xolair/asthma/capacity>