UNOVARTIS

Cosentyx Derm - SensoReady® 150 mg injection video - HCP

Prescribing information

Image



Image



SensoReady® 150 mg injection video

Cosentyx® (secukinumab) is indicated for the treatment of moderate to severe plaque psoriasis (PsO) in adults, children and adolescents from the age of 6 years who are candidates for systemic therapy; active psoriatic arthritis (PsA) in adult patients (alone or in combination with methotrexate [MTX]) when the response to previous disease-modifying anti-rheumatic drug therapy has been inadequate; active moderate to severe hidradenitis suppurativa (HS; acne inversa) in adults with an inadequate response to conventional systemic HS therapy.¹

Full indications for Cosentyx can be found here

Cosentyx is intended for use under the guidance and supervision of a physician experienced in the diagnosis and treatment of conditions for which Cosentyx is indicated.

This page contains a 5-minute video demonstrating how to prepare and deliver Cosentyx with the SensoReady $^{(\! 8\!)}$ pen.



For information on the Cosentyx UnoReady® 300 mg pen click here.





This video has been funded and developed by Novartis Pharmaceuticals UK Ltd.

Do not share screenshots or links to this website, video with patients, as this website is intended for healthcare professionals only. If you do wish to share a video with patients who have been prescribed Cosentyx, please share this link:

pro.novartis.com/uk-en/public/medicines/dermatology/cosentyx/patientresources/sensoready-injection-video

Click here for Cosentyx (secukinumab) prescribing information

Image

PsO dosing

Therapeutic indications¹

Cosentyx is indicated for the treatment of moderate to severe plaque psoriasis (PsO) in adults, children and adolescents from the age of 6 years who are candidates for systemic therapy; active psoriatic arthritis (PsA) in adult patients (alone or in combination with methotrexate [MTX]) when the response to previous disease-modifying anti-rheumatic drug therapy has been inadequate; active ankylosing spondylitis (AS) in adults who have responded inadequately to conventional therapy; active non-radiographic axial spondyloarthritis (nr-axSpA) with objective signs of inflammation as indicated by elevated C-reactive protein and/or magnetic resonance imaging evidence in adults who have responded inadequately to non-steroidal anti-inflammatory drugs; active moderate to severe hidradenitis suppurativa (HS; acne inversa) in adults with an inadequate response to conventional systemic HS therapy; active enthesitis-related arthritis (ERA) in patients 6 years and older (alone or in combination with MTX) whose disease has responded inadequately to, or who cannot tolerate, conventional therapy; active juvenile psoriatic arthritis (JPsA) in patients 6 years and older (alone or in combination with MTX) whose disease has responded inadequately to, or who cannot tolerate, conventional therapy.

AS, ankylosing spondylitis; CRP, C-reactive protein; DMARD, disease-modifying antirheumatic drug; ERA, enthesitis-related arthritis; HS, hidradenitis suppurativa; JPsA, juvenile psoriatic arthritis; MRI, magnetic resonance imaging; MTX, methotrexate; nr-axSpA, nonradiographic axial spondyloarthritis; NSAID, non-steroidal anti-inflammatory drug; PsA, psoriatic arthritis; PsO, plaque psoriasis.

References

- 1. Cosentyx® (secukinumab) GB Summary of Product Characteristics.
- 2. Cosentyx[®] (secukinumab) NI Summary of Product Characteristics.

UK | October 2024 | FA-11210365

Adverse events should be reported. Reporting forms and information can be found at <u>www.mhra.gov.uk/yellowcard</u>. Adverse events should also be reported to Novartis online through the pharmacovigilance intake (PVI) tool at <u>www.novartis.com/report</u>, or alternatively email <u>medinfo.uk@novartis.com</u> or call 01276 698370.

Source URL:

https://prod.pro.novartis.com/uk-en/medicines/dermatology/cosentyx/hcp-resources/sensor eady-injection-video